

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES


Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

$$\begin{array}{c} \bullet \\ \bullet \end{array} \quad \begin{array}{c} \text{---} \\ \text{---} \end{array} \quad \begin{array}{c} \bullet \\ \bullet \end{array}$$





Figure 2 hT1R2 and hT1R3 are expressed in human tongue epithelium. cDNA-specific amplification products can be amplified from cDNA prepared from resected human circumvallate papillae.

Figure 3 Human T1R2/T1R3 functions as a sweet taste receptor





Figure 4 T1R2 may control T1R2/T1R3 ligand specificity






Figure 5





Figure 6


Figure 7 Key ligand-binding residues of mGluR1 are conserved in T1R1



Figure 8 Human T1R1/T1R3 functions as an umami taste receptor



A.



B.




Figure 9 PDZIP facilitate the surface expression of human T1R2.

- A. Immunofluorescence staining of Myc-tagged hT1R2 indicates that PDZIP significantly increases the amount of human T1R2 protein on the plasma membrane.**
- B. FACS analysis data demonstrating the same result.
Myc-tagged human T1R2: Green line. Myc-tagged**
- C. human T1R2 with PDZIP: black line.**

Figure 10 Calcium-imaging data demonstrating hT1R2/hT1R3 responses to a number of sweet stimuli.




Figure 11




Figure 12





Figure 13






Figure 14




Figure 15




Figure 16 Lactisole inhibits the T1R2/T1R3 sweet and T1R1/T1R3 umami receptors and sweet and umami taste. (*Left panel*) responses of HEK- $G_{\alpha 15}$ cells transiently transfected with T1R1/T1R3 (*circles*) to 10 mM L-glutamate and HEK- $G_{\alpha 15}$ cells transiently transfected with T1R2/T1R3 (*squares*) to 150 mM sucrose in the presence of variable concentrations of lactisole are shown. (*Right panel*) fold increases in taste detection thresholds in the presence of 1 and 2 mM lactisole are shown for the sweet taste stimuli sucrose and D-tryptophan, the umami taste stimuli L-glutamate (MSG) and L-glutamate plus 0.2 mM IMP, and sodium chloride. Detection thresholds were determined following the method of Schiffman et al.